



Victorian Certificate of Education Year

CHEMISTRY

Written examination

Annotated 2019

DATA BOOK

by James Kennedy

Instructions

This data book is provided for your reference.
A question and answer book is provided with this data book.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

Table of contents

	Page
1. Periodic table of the elements	3
2. Electrochemical series	4
3. Chemical relationships	5
4. Physical constants and standard values	5
5. Unit conversions	6
6. Metric (including SI) prefixes	6
7. Acid-base indicators	6
8. Representations of organic molecules	7
9. Formulas of some fatty acids	7
10. Formulas of some biomolecules	8–9
11. Heats of combustion of common fuels	10
12. Heats of combustion of common blended fuels	10
13. Energy content of food groups	10
14. Characteristic ranges for infra-red absorption	11
15. ^{13}C NMR data	11
16. ^1H NMR data	12–13
17. 2-amino acids (α -amino acids)	14–15

ELECTROGEMATRY

**ALMOST ALWAYS
1+ AS AN ION
(except Li⁺, Na⁻)**

1. Periodic table of the elements

INCREASING COVALENT CHARGE

METALLIC CHARACTER
ATOMIC RADIUS

1 H	1.0 hydrogen	3 Li	4 Be	5 B	6 C	7 N-H	8 O-H	9 F-H	10 Ne
1.0		6.9 lithium	9.0 beryllium	10.8 boron	12.0 carbon	14.0 nitrogen	16.0 oxygen	19.0 fluorine	20.2 neon
hydrogen		lithium	beryllium	boron	carbon	nitrogen	oxygen	fluorine	neon
1 H	1.0 hydrogen	3 Li	4 Be	5 B	6 C	7 N-H	8 O-H	9 F-H	10 Ne
1.0		6.9 lithium	9.0 beryllium	10.8 boron	12.0 carbon	14.0 nitrogen	16.0 oxygen	19.0 fluorine	20.2 neon

1 H	1.0 hydrogen	3 Li	4 Be	5 B	6 C	7 N-H	8 O-H	9 F-H	10 Ne
1.0		6.9 lithium	9.0 beryllium	10.8 boron	12.0 carbon	14.0 nitrogen	16.0 oxygen	19.0 fluorine	20.2 neon
hydrogen		lithium	beryllium	boron	carbon	nitrogen	oxygen	fluorine	neon
1 H	1.0 hydrogen	3 Li	4 Be	5 B	6 C	7 N-H	8 O-H	9 F-H	10 Ne
1.0		6.9 lithium	9.0 beryllium	10.8 boron	12.0 carbon	14.0 nitrogen	16.0 oxygen	19.0 fluorine	20.2 neon

1 H	1.0 hydrogen	3 Li	4 Be	5 B	6 C	7 N-H	8 O-H	9 F-H	10 Ne
1.0		6.9 lithium	9.0 beryllium	10.8 boron	12.0 carbon	14.0 nitrogen	16.0 oxygen	19.0 fluorine	20.2 neon
hydrogen		lithium	beryllium	boron	carbon	nitrogen	oxygen	fluorine	neon
1 H	1.0 hydrogen	3 Li	4 Be	5 B	6 C	7 N-H	8 O-H	9 F-H	10 Ne
1.0		6.9 lithium	9.0 beryllium	10.8 boron	12.0 carbon	14.0 nitrogen	16.0 oxygen	19.0 fluorine	20.2 neon

ALWAYS 1+ AS IONS! ALWAYS SOLUBLE

CHEMISTRY DATA BOOK

TURN OVER

57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu
138.9 lanthanum	140.1 cerium	140.9 praseodymium	144.2 neodymium	145.0 promethium	150.4 samarium	152.0 europium	157.3 gadolinium	158.9 terbium	162.5 dysprosium	164.9 holmium	167.3 erbium	168.9 thulium	173.1 ytterbium	175.0 lutetium
lanthanum	cerium	praseodymium	neodymium	promethium	samarium	europium	gadolinium	terbium	dysprosium	holmium	erbium	thulium	ytterbium	lutetium
57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu
138.9 lanthanum	140.1 cerium	140.9 praseodymium	144.2 neodymium	145.0 promethium	150.4 samarium	152.0 europium	157.3 gadolinium	158.9 terbium	162.5 dysprosium	164.9 holmium	167.3 erbium	168.9 thulium	173.1 ytterbium	175.0 lutetium

89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr
(227) actinium	232.0 thorium	231.0 protactinium	238.0 uranium	(237) neptunium	(244) plutonium	(243) americium	(247) curium	(247) berkelium	(251) californium	(252) einsteinium	(257) fermium	(258) mendelevium	(259) nobelium	(262) lawrencium
actinium	thorium	protactinium	uranium	neptunium	plutonium	americium	curium	berkelium	californium	einsteinium	fermium	mendelevium	nobelium	lawrencium
89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr
(227) actinium	232.0 thorium	231.0 protactinium	238.0 uranium	(237) neptunium	(244) plutonium	(243) americium	(247) curium	(247) berkelium	(251) californium	(252) einsteinium	(257) fermium	(258) mendelevium	(259) nobelium	(262) lawrencium

The value in brackets indicates the mass number of the longest-lived isotope.

ALL ELEMENTS AFTER #2 ARE ARTIFICIAL →

CHEMISTRY DATA BOOK

CHEMISTRY DATA BOOK


CHEMISTRY DATA BOOK

2. Electrochemical series

ALL OF THESE HALF EQUATIONS SHOW REDUCTIONS

Reaction	Standard electrode potential (E°) in volts at 25 °C
OXIDANTS + $e^- \rightleftharpoons$ REDUCTANTS	
$F_2(g) + 2e^- \rightleftharpoons 2F^-(aq)$	+2.87
$H_2O_2(aq) + 2H^+(aq) + 2e^- \rightleftharpoons 2H_2O(l)$	+1.77
$Au^+(aq) + e^- \rightleftharpoons Au(s)$	+1.68
$Cl_2(g) + 2e^- \rightleftharpoons 2Cl^-(aq)$	+1.36
$O_2(g) + 4H^+(aq) + 4e^- \rightleftharpoons 2H_2O(l)$	+1.23
$Br_2(l) + 2e^- \rightleftharpoons 2Br^-(aq)$	+1.09
$Ag^+(aq) + e^- \rightleftharpoons Ag(s)$	+0.80
$Fe^{3+}(aq) + e^- \rightleftharpoons Fe^{2+}(aq)$	+0.77
$O_2(g) + 2H^+(aq) + 2e^- \rightleftharpoons H_2O_2(aq)$	+0.68
$I_2(s) + 2e^- \rightleftharpoons 2I^-(aq)$	+0.54
$O_2(g) + 2H_2O(l) + 4e^- \rightleftharpoons 4OH^-(aq)$	+0.40
$Cu^{2+}(aq) + 2e^- \rightleftharpoons Cu(s)$	+0.34
$Sn^{4+}(aq) + 2e^- \rightleftharpoons Sn^{2+}(aq)$	+0.15
$S(s) + 2H^+(aq) + 2e^- \rightleftharpoons H_2S(g)$	+0.14
$2H^+(aq) + 2e^- \rightleftharpoons H_2(g)$	0.00
$Pb^{2+}(aq) + 2e^- \rightleftharpoons Pb(s)$	-0.13
$Sn^{2+}(aq) + 2e^- \rightleftharpoons Sn(s)$	-0.14
$Ni^{2+}(aq) + 2e^- \rightleftharpoons Ni(s)$	-0.25
$Co^{2+}(aq) + 2e^- \rightleftharpoons Co(s)$	-0.28
$Cd^{2+}(aq) + 2e^- \rightleftharpoons Cd(s)$	-0.40
$Fe^{2+}(aq) + 2e^- \rightleftharpoons Fe(s)$	-0.44
$Zn^{2+}(aq) + 2e^- \rightleftharpoons Zn(s)$	-0.76
$2H_2O(l) + 2e^- \rightleftharpoons H_2(g) + 2OH^-(aq)$	-0.83
$Mn^{2+}(aq) + 2e^- \rightleftharpoons Mn(s)$	-1.18
$Al^{3+}(aq) + 3e^- \rightleftharpoons Al(s)$	-1.66
$Mg^{2+}(aq) + 2e^- \rightleftharpoons Mg(s)$	-2.37
$Na^+(aq) + e^- \rightleftharpoons Na(s)$	-2.71
$Ca^{2+}(aq) + 2e^- \rightleftharpoons Ca(s)$	-2.87
$K^+(aq) + e^- \rightleftharpoons K(s)$	-2.93
$Li^+(aq) + e^- \rightleftharpoons Li(s)$	-3.04

STRONGEST REDUCANT = MOST REACTIVE METAL ON THIS LIST



THESE VOLTAGES ALL CHANGE AT NON-STANDARD CONDITIONS

* STANDARD HYDROGEN ELECTRODE

FOR A SPONTANEOUS REACTION TO OCCUR:

REDUCTION CATHODE

OXIDATION ANODE

3. Chemical relationships

Name	Formula
number of moles of a substance	$n = \frac{m}{M}; n = cV; n = \frac{V}{V_m}$
universal gas equation	$pV = nRT$
calibration factor (CF) for bomb calorimetry	$CF = \frac{VIt}{\Delta T}$ <i>one term not 2</i> DON'T ADD 273 TO ΔT!
heat energy released in the combustion of a fuel (J)	$q = mc\Delta T$
enthalpy of combustion	$\Delta H = \frac{q}{n}$ (kJ) } CONVERT J TO KJ HERE BY ÷ 1000 IN CALCULATIONS
electric charge	$Q = It$
number of moles of electrons	$n(e^-) = \frac{Q}{F}$
% atom economy	$\frac{\text{molar mass of desired product}}{\text{molar mass of all reactants}} \times \frac{100}{1}$
% yield	$\frac{\text{actual yield}}{\text{theoretical yield}} \times \frac{100}{1}$

4. Physical constants and standard values

Name	Symbol	Value
Avogadro constant	N_A or L	$6.02 \times 10^{23} \text{ mol}^{-1}$
charge on one electron (elementary charge)	e	$-1.60 \times 10^{-19} \text{ C}$
Faraday constant	F	$96\,500 \text{ C mol}^{-1}$
molar gas constant	R	$8.31 \text{ J mol}^{-1} \text{ K}^{-1}$
molar volume of an ideal gas at SLC (25 °C and 100 kPa)	V_m	24.8 L mol^{-1}
specific heat capacity of water	c	4.18 kJ kg⁻¹ K⁻¹ or $4.18 \text{ J g}^{-1} \text{ K}^{-1}$
density of water at 25 °C	d	997 kg m^{-3} or 0.997 g mL^{-1}

$$n = \frac{N}{N_A}$$

$$C_1 V_1 = C_2 V_2$$

$$C_1 V_1 + C_2 V_2 = C_3 V_3$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$d = \frac{m}{V}$$

TURN OVER

RECENTLY CHANGED

5. Unit conversions

Measured value	Conversion
0 °C	273 K
100 kPa	750 mm Hg or 0.987 atm
1 litre (L)	1 dm ³ or 1×10^{-3} m ³ or 1×10^3 cm ³ or 1×10^3 mL $\Rightarrow 997\text{ g H}_2\text{O} @ 25^\circ\text{C}$

6. Metric (including SI) prefixes

Metric (including SI) prefixes	Scientific notation <i>IN CALCULATIONS, REPLACE PREFIXES WITH VALUES</i>	Multiplying factor
giga (G)	10^9	1 000 000 000
mega (M)	10^6	1 000 000
kilo (k)	10^3	1000
deci (d)	10^{-1}	0.1
centi (c)	10^{-2}	0.01
milli (m)	10^{-3}	0.001
micro (μ)	10^{-6}	0.000001
nano (n)	10^{-9}	0.000000001
pico (p)	10^{-12}	0.000000000001

pH FORMULAE

$$\text{pH} = -\log_{10} [\text{H}^+]$$

$$[\text{H}^+] = 10^{-\text{pH}}$$

$$[\text{H}^+] = \frac{10^{-14}}{[\text{OH}^-]}$$

ONLY TRUE IN AQUEOUS SOLUTIONS AT 25°C

$$[\text{OH}^-] = \frac{10^{-14}}{[\text{H}^+]}$$

$[\text{H}^+] = [\text{H}_3\text{O}^+]$ IN AQUEOUS SOLUTION

7. Acid-base indicators

Name	pH range	Colour change from lower pH to higher pH in range
thymol blue (1st change)	1.2–2.8	red \rightarrow yellow
methyl orange	3.1–4.4	red \rightarrow yellow
bromophenol blue	3.0–4.6	yellow \rightarrow blue
methyl red	4.4–6.2	red \rightarrow yellow
bromothymol blue	6.0–7.6	yellow \rightarrow blue
phenol red	6.8–8.4	yellow \rightarrow red
thymol blue (2nd change)	8.0–9.6	yellow \rightarrow blue
phenolphthalein	8.3–10.0	colourless \rightarrow pink 

IN PURE WATER,
 $[\text{H}^+] = [\text{OH}^-]$
 EVEN IF THE pH ISN'T 7

8. Representations of organic molecules

The following table shows different representations of organic molecules, using butanoic acid as an example.

Formula	Representation
molecular formula	$\text{C}_4\text{H}_8\text{O}_2$
structural formula	<p>YOU MUST DRAW ALL ATOMS & ALL BONDS.</p>
semi-structural (condensed) formula	$\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$ or $\text{CH}_3(\text{CH}_2)_2\text{COOH}$
skeletal structure	<p>CORNERS & ENDS WITHOUT A LETTER ARE CARBON ATOMS.</p>

→ ALL CONTAIN CARBOXYL GROUP $(-\text{C}^{\text{II}}\text{O}-\text{H})$

9. Formulas of some fatty acids → ALL ARE SOLID AT 25°C

→ Name	Formula	$N(\text{C}=\text{C})$	Semi-structural formula
lauric acid	$\text{C}_{12}\text{H}_{24}\text{COOH}$	0	$\text{CH}_3(\text{CH}_2)_{10}\text{COOH}$
myristic acid	$\text{C}_{14}\text{H}_{28}\text{COOH}$	0	$\text{CH}_3(\text{CH}_2)_{12}\text{COOH}$
palmitic acid	$\text{C}_{16}\text{H}_{32}\text{COOH}$	0	$\text{CH}_3(\text{CH}_2)_{14}\text{COOH}$
palmitoleic acid	$\text{C}_{16}\text{H}_{30}\text{COOH}$	1	$\text{CH}_3(\text{CH}_2)_4\text{CH}_2\text{CH}=\text{CHCH}_2(\text{CH}_2)_5\text{CH}_2\text{COOH}$
stearic acid	$\text{C}_{18}\text{H}_{36}\text{COOH}$	0	$\text{CH}_3(\text{CH}_2)_{16}\text{COOH}$
oleic acid	$\text{C}_{18}\text{H}_{34}\text{COOH}$	1	$\text{CH}_3(\text{CH}_2)_7\text{CH}=\text{CH}(\text{CH}_2)_7\text{COOH}$
linoleic acid	$\text{C}_{18}\text{H}_{32}\text{COOH}$	2	$\text{CH}_3(\text{CH}_2)_4(\text{CH}=\text{CHCH}_2)_2(\text{CH}_2)_6\text{COOH}$
linolenic acid	$\text{C}_{18}\text{H}_{30}\text{COOH}$	3	$\text{CH}_3\text{CH}_2(\text{CH}=\text{CHCH}_2)_3(\text{CH}_2)_6\text{COOH}$
arachidic acid	$\text{C}_{20}\text{H}_{38}\text{COOH}$	0	$\text{CH}_3(\text{CH}_2)_{17}\text{CH}_2\text{COOH}$
arachidonic acid	$\text{C}_{20}\text{H}_{36}\text{COOH}$	4	$\text{CH}_3(\text{CH}_2)_4(\text{CH}=\text{CHCH}_2)_3\text{CH}=\text{CH}(\text{CH}_2)_3\text{COOH}$

TO FIND $N(\text{C}=\text{C})$ IN A FATTY ACID TAIL, USE:

$$N(\text{C}=\text{C}) = C - \frac{H-1}{2}$$



(e.g.) Arachidonic Acid $\text{C}_{20}\text{H}_{36}\text{COOH}$

$$N(\text{C}=\text{C}) = 20 - \frac{36-1}{2} = 4$$

TURN OVER


10. Formulas of some biomolecules

NON-POLAR




vitamin D₂ (ergocalciferol) ← NOT STRUCTURAL ISOMERS → vitamin D₃ (cholecalciferol)

POLAR




vitamin C (ascorbic acid)




glycerol

MONOSACCHARIDES

$C_6H_{12}O_6$
 $M_r = 180$




α -glucose



β -fructose (HAS FIVE SIDES)


DISACCHARIDES

$C_{12}H_{22}O_{11}$
 $M_r = 342$





sucrose


ETHER / GLYCOSIDIC LINK
FORMS DURING A CONDENSATION
REACTION
releasing H_2O as
a product



α -lactose



amylopectin (starch)



amylose (starch)

TURN OVER

11. Heats of combustion of common fuels

The heats of combustion in the following table are calculated at SLC (25 °C and 100 kPa) with combustion products being CO₂ and H₂O. Heat of combustion may be defined as the heat energy released when a specified amount of a substance burns completely in oxygen and is, therefore, reported as a positive value, indicating a magnitude. Enthalpy of combustion, ΔH , for the substances in this table would be reported as negative values, indicating the exothermic nature of the combustion reaction.

VERY USEFUL INFO

Fuel	Formula	State	Heat of combustion (kJ g ⁻¹)	Molar heat of combustion (kJ mol ⁻¹)
hydrogen	H ₂	gas	141	282
methane	CH ₄	gas	55.6	890
ethane	C ₂ H ₆	gas	51.9	1560
propane	C ₃ H ₈	gas	50.5	2220
butane	C ₄ H ₁₀	gas	49.7	2880
octane	C ₈ H ₁₈	liquid	47.9	5460
ethyne (acetylene)	C ₂ H ₂	gas	49.9	1300
methanol	CH ₃ OH	liquid	22.7	726
ethanol	C ₂ H ₅ OH	liquid	29.6	1360

NOTHING AQUEOUS COMBUSTS (ACCORDING TO VCAA)

12. Heats of combustion of common blended fuels

Blended fuels are mixtures of compounds with different mixture ratios and, hence, determination of a generic molar enthalpy of combustion is not realistic. The values provided in the following table are typical values for heats of combustion at SLC (25 °C and 100 kPa) with combustion products being CO₂ and H₂O. Values for heats of combustion will vary depending on the source and composition of the fuel.

Fuel ALL MIXTURES ALL FOSSIL FUELS	State	Heat of combustion (kJ g ⁻¹)
kerosene	liquid	46.2
diesel	liquid	45.0
natural gas	gas	54.0

BALANCE COMBUSTION EQUATIONS:

C, H, O.

HALVES & TOP-HEAVY FRACTIONS
ARE RECOMMENDED! (eg $\underline{\underline{2\frac{1}{2}}}, \underline{\underline{\frac{5}{2}}}$)

13. Energy content of food groups

Food	Heat of combustion (kJ g ⁻¹)	DIGESTED	BY (ENZYMES)
fats and oils	37	SMALL INTESTINE	PANCREATIC LIPASE
protein	17	STOMACH SMALL + LARGE INT.	PEPSIN TRYPsin, etc.
carbohydrate	16	SALIVA AND SMALL INTESTINE.	AMYLASE.

ALWAYS COPY BOND NAMES EXACTLY

14. Characteristic ranges for infra-red absorption

Bond	Wave number (cm ⁻¹)	Bond	Wave number (cm ⁻¹)
C—Cl (chloroalkanes)	600–800	C=O (ketones)	1680–1850
C—O (alcohols, esters, ethers)	1050–1410	C=O (esters)	1720–1840
C=C (alkenes)	1620–1680	C—H (alkanes, alkenes, arenes)	2850–3090
C=O (amides)	1630–1680	O—H (acids)	2500–3500
C=O (aldehydes)	1660–1745	O—H (alcohols)	3200–3600
C=O (acids)	1680–1740	N—H (amines and amides)	3300–3500

15. ¹³C NMR dataTypical ¹³C shift values relative to TMS = 0

These can differ slightly in different solvents.



Type of carbon	Chemical shift (ppm)	THERE'S NO SPLITTING IN ¹³ C NMR
R—CH ₃ METHYL	8–25	
R—CH ₂ —R STRAIGHT	20–45	
R ₃ —CH BRANCHED	40–60	
R ₄ —C DOUBLE-BRANCHED	36–45	
R—CH ₂ —X 1-HALO-	15–80	
R ₃ C—NH ₂ , R ₃ C—NR AMINE	35–70	
R—CH ₂ —OH PRIMARY ALCOHOL	50–90	
RC≡CR ALKYNE	75–95	
R ₂ C=CR ₂ ALKENE	110–150	
RCOOH CARBOXYL	160–185	
ESTER	165–175	↗
ALDEHYDE	190–200	
R ₂ C=O KETONE	205–220	





16. ^1H NMR data

Typical proton shift values relative to TMS = 0

These can differ slightly in different solvents. The shift refers to the proton environment that is indicated in bold letters in the formula.

Type of proton	Chemical shift (ppm)
$\text{R}-\text{CH}_3$ METHYL	0.9–1.0
$\text{R}-\text{CH}_2-\text{R}$ STRAIGHT	1.3–1.4
$\text{RCH}=\text{CH}-\text{CH}_3$ -2-ENE	1.6–1.9
R_3-CH BRANCHED	1.5
 or	2.0
 KETONE (-2-ONE)	2.1–2.7
$\text{R}-\text{CH}_2-\text{X}$ ($\text{X} = \text{F}, \text{Cl}, \text{Br}$ or I) 1-HALO-	3.0–4.5
 PRIMARY ALCOHOL → SECONDARY ALCOHOL	3.3–4.5
 SECONDARY AMIDE	3.2
$\text{R}-\text{O}-\text{CH}_3$ or $\text{R}-\text{O}-\text{CH}_2\text{R}$	3.3–3.7
 PHENYL ETHANATE MOLECULE	2.3
 ESTER	3.7–4.8
$\text{R}-\text{O}-\text{H}$ ALWAYS A SINGLET ALCOHOL	1–6 (varies considerably under different conditions)
$\text{R}-\text{NH}_2$ PRIMARY AMINE	1–5
$\text{RHC}=\text{CHR}$ -1-ENE	4.5–7.0
 PHENOL	4.0–12.0

singlet doublet triplet quartet ... multiplet
 1 2 3 4 ... n

Type of proton	Chemical shift (ppm)
 BENZENE	6.9–9.0 ⚡
 SECONDARY AMIDE	8.1
 ALDEHYDE	9.4–10.0 ⚡
 CARBOXYLIC ACID ← ALWAYS A SINGLET	9.0–13.0

ALWAYS SOLUBLE IN WATER



X^- (EXCEPT FOR SILVER, MERCURY & LEAD SALTS)

SO_4^{2-} (EXCEPT FOR BARIUM, CALCIUM & LEAD SALTS)

NEVER SOLUBLE IN WATER

S^{2-} (EXCEPT K^+ Na^+ NH_4^+)

CO_3^{2-} (EXCEPT K^+ Na^+ NH_4^+)

PO_4^{3-} (EXCEPT K^+ Na^+ NH_4^+)

OH^- (EXCEPT K^+ Na^+ NH_4^+)

NON-POLAR
 POLAR
 ACIDIC
 BASIC

17. 2-amino acids (α -amino acids)

The table below provides simplified structures to enable the drawing of zwitterions, the identification of products of protein hydrolysis and the drawing of structures involving condensation polymerisation of amino acid monomers.

Name	Symbol	Structure
alanine	Ala	$\text{H}_2\text{N}—\text{CH}(\text{CH}_3)—\text{COOH}$
arginine	Arg	$\text{H}_2\text{N}—\text{CH}(\text{CH}_2\text{CH}_2\text{CH}_2\text{NH})—\text{C}(=\text{O})\text{NH}_2$
asparagine	Asn	$\text{H}_2\text{N}—\text{CH}(\text{CH}_2\text{C}(=\text{O})\text{NH}_2)—\text{COOH}$ <p style="color:red; margin-left: 20px;">AMIDE GROUP IS POLAR NOT BASIC</p>
aspartic acid	Asp	$\text{H}_2\text{N}—\text{CH}(\text{CH}_2\text{COOH})—\text{COOH}$
cysteine	Cys	$\text{H}_2\text{N}—\text{CH}(\text{CH}_2\text{SH})—\text{COOH}$ <p style="color:red; margin-left: 20px;">SULFUR-CONTAINING</p>
glutamic acid	Glu	$\text{H}_2\text{N}—\text{CH}(\text{CH}_2\text{CH}_2\text{COOH})—\text{COOH}$
glutamine	Gln	$\text{H}_2\text{N}—\text{CH}(\text{CH}_2\text{CH}_2\text{C}(=\text{O})\text{NH}_2)—\text{COOH}$ <p style="color:red; margin-left: 20px;">AMIDE GROUP IS POLAR NOT BASIC</p>
glycine	Gly	$\text{H}_2\text{N}—\text{CH}_2—\text{COOH}$
histidine	His	$\text{H}_2\text{N}—\text{CH}(\text{CH}_2\text{NHC}_6\text{H}_4\text{CH}_2\text{NH}_2)—\text{COOH}$
isoleucine	Ile	$\text{H}_2\text{N}—\text{CH}(\text{CH}_3\text{CH}_2\text{CH}_3)—\text{COOH}$

Name	Symbol	Structure
leucine	Leu	$\begin{array}{c} \text{CH}_3-\text{CH}-\text{CH}_3 \\ \\ \text{CH}_2 \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
lysine	Lys	$\begin{array}{c} \text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}_2 \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
methionine	Met	$\begin{array}{c} \text{CH}_2-\text{CH}_2-\text{S}-\text{CH}_3 \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
phenylalanine	Phe	$\begin{array}{c} \text{CH}_2-\text{C}_6\text{H}_5 \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
proline	Pro	
serine	Ser	$\begin{array}{c} \text{CH}_2-\text{OH} \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
threonine	Thr	$\begin{array}{c} \text{CH}_3-\text{CH}-\text{OH} \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
tryptophan	Trp	$\begin{array}{c} \text{CH}_2-\text{C}_6\text{H}_4-\text{NH} \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$ <p style="color:red; margin-left: 20px;">CONSIDERED NON-POLAR DESPITE THE "NH".</p>
tyrosine	Tyr	$\begin{array}{c} \text{CH}_2-\text{C}_6\text{H}_4-\text{OH} \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
valine	Val	$\begin{array}{c} \text{CH}_3-\text{CH}-\text{CH}_3 \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$